



# Commentary: Maternal immune activation evoked by polyinosinic: polycytidylic acid does not evoke microglial cell activation in the embryo

Hans-Gert Bernstein<sup>1\*</sup>, Yael Piontkewitz<sup>2</sup> and Gerburg Keilhoff<sup>3</sup>

<sup>1</sup> Department of Psychiatry, University of Magdeburg, Magdeburg, Germany, <sup>2</sup> George S. Wise Faculty of Life Sciences, Strauss Center for Computational Neuroimaging, Tel Aviv University, Tel Aviv, Israel, <sup>3</sup> Department of Biochemistry and Cell Biology, University of Magdeburg, Magdeburg, Germany

**Keywords:** maternal immune activation, Poly I:C, Microglia, astrocytes, offspring

## A commentary on

### Maternal immune activation evoked by polyinosinic: polycytidylic acid does not evoke microglial cell activation in the embryo

by Smolders, S., Smolders, S. M., Swinnen, N., Gärtner, A., Rigo, J. M., Legendre, P., et al. (2015). *Front. Cell. Neurosci.* 9:301. doi: 10.3389/fncel.2015.00301

## OPEN ACCESS

### Edited by:

Takahiro A. Kato,  
Kyushu University, Japan

### Reviewed by:

Manabu Makinodan,  
Nara Medical University, Japan

### \*Correspondence:

Hans-Gert Bernstein  
hans-gert.bernstein@med.ovgu.de

**Received:** 15 December 2015

**Accepted:** 05 February 2016

**Published:** 23 February 2016

### Citation:

Bernstein H-G, Piontkewitz Y and Keilhoff G (2016) Commentary: Maternal immune activation evoked by polyinosinic: polycytidylic acid does not evoke microglial cell activation in the embryo. *Front. Cell. Neurosci.* 10:41. doi: 10.3389/fncel.2016.00041

Immune-related abnormalities, which probably result from maternal infections during pregnancy, can be found in patients with schizophrenia and other mental disorders. In the endeavor to simulate this environmental schizophrenia risk in animal models, maternal immune activation (MIA) by infectious agents has been introduced (Zuckerman et al., 2003; Meyer et al., 2005). In a majority of publications on MIA, either lipopolysaccharide (LPS), a cell wall component of Gram-negative bacteria, or the viral mimetic polyriboinosinic-polyribocytidylic acid (poly I:C), are administered to pregnant mouse or rat dams. These maternal immune challenges are considered as suitable schizophrenia paradigms, since they induce characteristic (and often similar) anatomical, cellular, neurochemical, and behavioral alterations in the offspring, which are of relevance for schizophrenia (Meyer et al., 2009 and many others). MIA with either LPS or poly I:C generates a broad immune-inflammatory response in the developing CNS of the offspring (for recent reviews see Dean et al., 2015; Giovanoli et al., 2015a; Smolders et al., 2015). However, the action of both agents might differ with regard to one remarkable aspect: while LPS activates microglia *in vivo* and *in vitro* (Roumier et al., 2008; Cunningham, 2013; Dean et al., 2015; Zager et al., 2015 and others), poly I:C possibly does not (no activation: Olson and Miller, 2004; Piontkewitz et al., 2012; Giovanoli et al., 2015a,b; Smolders et al., 2015; activation: Patro et al., 2010; Juckel et al., 2011; Missault et al., 2014; Van den Eynde et al., 2014; Zhu et al., 2014). In an elegant set of *in vivo* and *in vitro* experiments Smolders et al. (2015) examined the effect of LPS and poly I:C under identical conditions. In particular, they investigated whether embryonic microglia can be directly activated by incubating mouse brain slices from embryonic day 15.5 with either saline, poly I:C, IL-6, or LPS. They found that LPS, contrary to poly I:C or IL-6, activates microglia to “a detrimental activation state.” When discussing possible pathophysiologic consequences of poly I:C’s failure to activate microglia, Smolders et al. (2015) come to three conclusions: (i) It is unlikely that embryonic microglia dysfunction is the main mechanism that induces developmental abnormalities, (ii) poly I:C might evoke developmental deficits by directly acting on neuronal development, and

(iii) it cannot be excluded that poly I:C effectuates an embryonic microglia priming, which results in an exaggerated response of microglia. Apart from “priming of microglia by poly I:C” being an exciting idea, there is yet little experimental evidence in favor of the existence of such a mechanism (perhaps via changes in the microglial kynurenine pathway? Giovanoli et al., 2015b). Hence, we would like to concentrate on the first two assumptions. Let's begin with the second one: is it conceivable (and plausible) that poly I:C induces developmental deficits by directly acting on neuronal development? In our opinion the answer is yes. Poly I:C is a strong agonist of Toll-like receptor 3 (TLR3). This receptor is already expressed in very immature neurons (Shi et al., 2013), and becomes up-regulated in a subpopulation of neurons after the injection of poly I:C (Deleidi et al., 2010). Moreover, poly I:C was found to depress embryonic neuronal stem cell division and population of the superficial layers of the neocortex by neurons, which was not the case with TLR3 deficient animals (De Miranda et al., 2010). And lastly, it has been shown that poly I:C treatment of pregnant rat dams leads to an impaired postnatal neurogenesis, but not disturbed microgliogenesis (Piontkewitz et al., 2012), as well as to an impaired adult neurogenesis (Zhang and van Praag, 2015), in the hippocampus of the offspring. Thus, poly I:C might well exert direct influence on neuronal development as proposed. However, this interaction can hardly explain the poly I:C induced cerebral immune-inflammatory response in the offspring. And this brings us back to the initial statement of Smolders and co-workers, namely, that microglia cannot be a main player in poly I:C induced developmental deficits. Assuming that this supposition is correct (some aforementioned *in vitro* and *in vivo* studies argue against this conjecture) one has to ask which brain tissue component then is to blame for the observed alterations, especially for the immune response? A “hot candidate” for this is astroglia. Astrocytes are abundantly populated with TLR3 (Farina et al., 2005, 2007; Park et al., 2006; Ibi et al., 2013; Ibi and Yamada, 2015 and others), become strongly activated

after poly I:C and, most importantly in this context, are able to secrete the whole battery of pro-inflammatory and anti-inflammatory cytokines, which are typically found after MIA with poly I:C (as reviewed by Ibi and Yamada, 2015). Moreover, when cultured neurons were incubated with the conditioned medium of poly I:C treated astrocytes, neurite development was found to be disturbed. This effect is mediated by an interferon-induced transmembrane protein 3, which is synthesized by, and released into the medium from, astrocytes after poly I:C treatment (Ibi et al., 2013). Analysis of conditioned media of astrocytes after poly I:C treatment subsequently revealed the presence of a further protein, matrix metalloproteinase 3, which also contributes to the observed impairment of neurite outgrowth and spine formation of cultured neurons. Of note, this protein is expressed in, and released from, astrocytes but not microglia (Yamada et al., 2014). Moreover, strong astroglial activation may be detected in postnatal hippocampus of the offspring after mid-gestational poly I:C MIA using GFAP immunolabeling (Ratnayake et al., 2012). Interestingly, Ibi and Yamada (2015) claim that poly I:C activates TLR3 in astrocytes of the brain parenchyma or BBB, thus pointing to a possible role of activated astroglia in impaired vascularization. Indeed, TLR3 activation has a pronounced anti-angiogenic effect (Grellet et al., 2013), but it is yet not fully clear, if astroglia is implicated in this process. In any case, impaired vascularization was found by reduced RECA-1 immunohistochemistry in postnatal rat hippocampi after MIA by poly I:C treatment (Piontkewitz et al., 2012). In sum, there are good reasons to consider astroglia as a major player in brain pathology of the offspring (including immune-inflammatory response) after maternal exposure to poly I:C.

## AUTHOR CONTRIBUTIONS

All authors listed, have made substantial, direct and intellectual contribution to the work, and approved it for publication.

## REFERENCES

- Cunningham, C. (2013). Microglia and neurodegeneration: the role of systemic inflammation. *Glia* 61, 71–90. doi: 10.1002/glia.22350
- Dean, J. M., Shi, Z., Fleiss, B., Gunn, K. C., Groenendaal, F., van Bel, F., et al. (2015). A critical review of models of perinatal infection. *Dev. Neurosci.* 37, 289–304. doi: 10.1159/000370309
- Deleidi, M., Hallett, P. J., Koprich, J. B., Chung, C. Y., and Isacson, O. (2010). The Toll-like receptor-3 agonist polyinosinic:polycytidylic acid triggers nigrostriatal dopaminergic degeneration. *J. Neurosci.* 30, 16091–16101. doi: 10.1523/JNEUROSCI.2400-10.2010
- De Miranda, J., Yaddanapudi, K., Hornig, M., Villar, G., Serge, R., and Lipkin, W. I. (2010). Induction of Toll-like receptor 3-mediated immunity during gestation inhibits cortical neurogenesis and causes behavioral disturbances. *MBio* 1, e00176–e00110. doi: 10.1128/mbio.00176-10
- Farina, C., Krumbholz, M., Giese, T., Hartmann, G., Aloisi, F., and Meinl, E. (2005). Preferential expression and function of Toll-like receptor 3 in human astrocytes. *J. Neuroimmunol.* 159, 12–19. doi: 10.1016/j.jneuroim.2004.09.009
- Farina, C., Aloisi, F., and Meinl, E. (2007). Astrocytes are active players in cerebral innate immunity. *Trends Immunol.* 28, 138–145. doi: 10.1016/j.it.2007.01.005
- Giovanoli, S., Notter, T., Richetto, J., Labouesse, M. A., Vuillermot, S., Riva, M. A., et al. (2015a). Late prenatal immune activation causes hippocampal deficits in the absence of persistent inflammation across aging. *J. Neuroinflammation* 12, 221. doi: 10.1186/s12974-015-0437-y
- Giovanoli, S., Weber-Stadlbauer, U., Schedlowski, M., Meyer, U., and Engler, H. (2015b). Prenatal immune activation causes hippocampal synaptic deficits in the absence of overt microglia anomalies. *Brain. Behav. Immunity*. doi: 10.1016/j.bbi.2015.09.015. [Epub ahead of print].
- Grellet, A., Cras, A., Balitrand, N., Delmau, C., Lecourt, S., Lepelletier, Y., et al. (2013). Toll-like receptor 3 regulates cord blood-derived endothelial cell function *in vitro* and *in vivo*. *Angiogenesis* 16, 821–836. doi: 10.1007/s10456-013-9358-5
- Ibi, D., Nagai, T., Nakajima, A., Mizoguchi, H., Kawase, T., Tsuboi, D., et al. (2013). Astroglial IFITM3 mediates neuronal impairments following neonatal immune challenge in mice. *Glia* 61, 679–693. doi: 10.1002/glia.22461
- Ibi, D., and Yamada, K. (2015). Therapeutic targets for neurodevelopmental disorders emerging from animal models with perinatal immune activation. *Int. J. Mol. Sci.* 16, 28218–28229. doi: 10.3390/ijms161226092
- Juckel, G., Manitz, M. P., Brüne, M., Friebe, A., Heneka, M. T., and Wolf, R. J. (2011). Microglial activation in a neuroinflammatory animal model of schizophrenia—a pilot study. *Schizophr. Res.* 131, 96–100. doi: 10.1016/j.schres.2011.06.018
- Meyer, U., Feldon, J., Schedlowski, M., and Yee, B. K. (2005). Towards an immuno-precipitated neurodevelopmental animal model of schizophrenia.

- Neurosci. Biobehav. Rev.* 29, 913–947. doi: 10.1016/j.neubiorev.2004.10.012
- Meyer, U., Feldon, J., and Fatemi, S. H. (2009). *In-vivo* rodent models for the experimental investigation of prenatal immune activation effects in neurodevelopmental brain disorders. *Neurosci Biobehav. Rev.* 33, 1061–1079. doi: 10.1016/j.neubiorev.2009.05.001
- Missault, S., Van den Eynde, K., Vanden Berghe, W., Fransen, E., Weeren, A., Timmermans, J. P., et al. (2014). The risk for behavioural deficits is determined by the maternal immune response to prenatal immune challenge in a neurodevelopmental model. *Brain Behav Immun.* 42, 138–146. doi: 10.1016/j.bbi.2014.06.013
- Olson, J. K., and Miller, S. D. (2004). Microglia initiate central nervous system innate and adaptive immune responses through multiple TLRs. *J. Immunol.* 173, 3916–3924. doi: 10.4049/jimmunol.173.6.3916
- Park, C., Lee, S., Cho, I. H., Lee, H. K., Kim, D., Choi, S. Y., et al. (2006). TLR3-mediated signal induces proinflammatory cytokine and chemokine gene expression in astrocytes: differential signaling mechanisms of TLR3-induced IP-10 and IL-8 gene expression. *Glia* 53, 248–256. doi: 10.1002/glia.20278
- Patro, I. K., Amit, S. M., Bhumika, S., and Patro, N. (2010). Poly I:C induced microglial activation impairs motor activity in adult rats. *Indian J. Exp. Biol.* 48, 104–109.
- Piontkewitz, Y., Bernstein, H. G., Dobrowolny, H., Bogerts, B., Weiner, I., and Keilhoff, G. (2012). Effects of risperidone treatment in adolescence on hippocampal neurogenesis, parvalbumin expression, and vascularization following prenatal immune activation in rats. *Brain Behav. Immun.* 26, 353–363. doi: 10.1016/j.bbi.2011.11.004
- Ratnayake, U., Quinn, T. A., Castillo-Melendez, M., Dickinson, H., and Walker, D. W. (2012). Behaviour and hippocampus-specific changes in spiny mouse neonates after treatment of the mother with the viral-mimetic Poly I:C at mid-pregnancy. *Brain Behav. Immun.* 26, 1288–1299. doi: 10.1016/j.bbi.2012.08.011
- Roumier, A., Pascual, O., Béchade, C., Wakselman, S., Poncer, J. C., Réal, E., et al. (2008). Prenatal activation of microglia induces delayed impairment of glutamatergic synaptic function. *PLoS ONE* 3:e2595. doi: 10.1371/journal.pone.0002595
- Shi, H., Gabarin, N., Hickey, E., and Askalan, R. (2013). TLR-3 receptor activation protects the very immature brain from ischemic injury. *J. Neuroinflammation* 10:104. doi: 10.1186/1742-2094-10-104
- Smolders, S., Smolders, S. M., Swinnen, N., Gärtner, A., Rigo, J. M., Legendre, P., et al. (2015). Maternal immune activation evoked by polyinosinic: polycytidylic acid does not evoke microglial cell activation in the embryo. *Front. Cell. Neurosci.* 9:301. doi: 10.3389/fncel.2015.00301
- Van den Eynde, K., Missault, S., Fransen, E., Raeymaekers, L., Willems, R., Drinkenburg, W., et al. (2014). Hypolocomotive behaviour associated with increased microglia in a prenatal immune activation model with relevance to schizophrenia. *Behav. Brain Res.* 258, 179–186. doi: 10.1016/j.bbr.2013.10.005
- Yamada, S., Nagai, T., Nakai, T., Ibi, D., Nakajima, A., and Yamada, K. (2014). Matrix metalloproteinase-3 is a possible mediator of neurodevelopmental impairment due to polyI:C-induced innate immune activation of astrocytes. *Brain Behav. Immun.* 38, 272–282. doi: 10.1016/j.bbi.2014.02.014
- Zager, A., Peron, J. P., Mennezier, G., Rodrigues, S. C., Aloia, T. P., and Palermo-Neto, J. (2015). Maternal immune activation in late gestation increases neuroinflammation and aggravates experimental autoimmune encephalomyelitis in the offspring. *Brain Behav. Immun.* 43, 159–171. doi: 10.1016/j.bbi.2014.07.021
- Zhang, Z., and van Praag, H. (2015). Maternal immune activation differentially impacts mature and adult-born hippocampal neurons in male mice. *Brain Behav. Immun.* 45, 60–70. doi: 10.1016/j.bbi.2014.10.010
- Zhu, F., Zheng, Y., Liu, Y., Zhang, X., and Zhao, J. (2014). Minocycline alleviates behavioral deficits and inhibits microglial activation in the offspring of pregnant mice after administration of polyriboinosinic-polyribocytidylic acid. *Psychiatry Res.* 219, 680–686. doi: 10.1016/j.psychres.2014.06.046
- Zuckerman, L., Rehavi, M., Nachman, R., and Weiner, I. (2003). Immune activation during pregnancy in rats leads to a postpubertal emergence of disrupted latent inhibition, dopaminergic hyperfunction, and altered limbic morphology in the offspring: a novel neurodevelopmental model of schizophrenia. *Neuropsychopharmacology* 28, 1778–1789. doi: 10.1038/sj.npp.1300248

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Bernstein, Piontkewitz and Keilhoff. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.